

**Supplementary Table 2.** Patient Global Impression (PGI) at the end of treatment

Variable	Vibegron 50 mg (n = 85)	Placebo (n = 95)	P-value
Satisfied <sup>c)</sup>			
No. (%)	75 (88.24)	67 (70.53)	0.0037 <sup>a)</sup>
Difference in responder rates (%)	17.71	-	
95% CI	6.26–29.15	-	
Very satisfie <sup>d)</sup>			<0.0001 <sup>a)</sup>
No. (%)	58 (68.24)	34 (35.79)	
Difference in responder rates (%)	32.45	-	
95% CI	18.63–46.26	-	
PGI score			0.0004 <sup>b)</sup>
1. Very much improved	23 (27.06)	15 (15.79)	
2. Much improved	35 (41.18)	19 (20.00)	
3. Improved	17 (20.00)	33 (34.74)	
4. No change	10 (11.76)	25 (26.32)	
5. Minimally worse	0 (0.0)	2 (2.11)	
6. Much worse	0 (0.0)	1 (1.05)	
7. Very much worse	0 (0.0)	0 (0.00)	

Values are presented as number (%).

CI, confidence interval.

<sup>a)</sup>Chi-square test. <sup>b)</sup>Fisher exact test. <sup>c)</sup>PGI ≤ 3. <sup>d)</sup>PGI ≤ 2.